

UNCLASSIFIED

AD 297 337

*Reproduced
by the*

**ARMED SERVICES TECHNICAL INFORMATION AGENCY
ARLINGTON HALL STATION
ARLINGTON 12, VIRGINIA**



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

63-2-5

NEL REPORT 1155

CATALOGED ATIA
As AL NO. 100-1007

297 337

CORRELATION BETWEEN THE MINIMUM VIRTUAL E-LAYER HEIGHT $h'E$ AND THE 100-mb CIRCULATION

V. R. Noonkester

Research Report 1155

22 January 1963

U. S. NAVY ELECTRONICS LABORATORY, SAN DIEGO, CALIFORNIA • A BUREAU OF SHIPS LABORATORY

18111
MAR 4 1963

THE PROBLEM

Study vlf and lf electromagnetic propagation mechanisms. Study the interrelationships of phenomena within the upper stratosphere and ionosphere which affect vlf and lf electromagnetic propagation and the extent to which variations of these phenomena may influence radio propagation. The study reported here examined the correlation between the minimum virtual E -layer height $h'E$ at Washington, D.C., and the stratospheric circulation over the eastern United States as a function of frequency and scale size of the circulation.

RESULTS

The minimum virtual E -layer height was found to be correlated with a particular type of large-scale circulation at the 100-mb level which has a period of 6.3 days. One of the ten circulation indices used to describe the 100-mb circulation was found to have predominant periods of 22 and 12 days. The predominant 12-day period appears to be controlling the correlation between the 100-mb circulation and $h'E$ at the 6.3-day period.

RECOMMENDATION

Correlate other ionospheric parameters with the circulation over larger regions of the world and at other pressure levels in the stratosphere or troposphere. Examine the physics of the relationship between $h'E$ and

the stratospheric circulation particularly with respect to the preference for $h'E$ to be related with certain circulation patterns which occur at certain periods.

ADMINISTRATIVE INFORMATION

Work was performed under SR 008 01 01, Task 7039 (NEL M2-3) and SF 011 05 11, Task 0558 (NEL M2-5) by members of the Radio Physics Division. The report covers intermittent work from 1 January 1958 to October 1962 and was approved for publication 22 January 1963.

The author expresses his appreciation to W. F. Moler and Dr. E. E. Gossard for their many suggestions and continued encouragements made during the progress of this study. Special thanks are extended to F. A. Sabransky who skillfully programmed the digital computer for the extensive calculations required and patiently altered these programs at my request. Without the careful work of R. O. Niemi in processing the data, considerable time would have been wasted. The careful preparation of the manuscript by Mrs. Eileen Jacobson is appreciated.

CONTENTS

INTRODUCTION...	page 5
DEFINITION OF VARIABLES...	5
CIRCULATION INDICES...	8
POWER SPECTRUM...	13
CORRELATION BETWEEN \bar{h}' AND THE 100-mb CIRCULATION...	18
SYNOPTIC EXAMPLE...	23
COHERENCE BETWEEN THE CIRCULATION INDICES...	26
CONCLUSIONS...	27
RECOMMENDATION...	27
REFERENCES...	28

ILLUSTRATIONS

- 1 Map of seven stations used to describe the 100-mb
circulation... page 6
- 2 Response curve of mathematical filter... 7
- 3 Distribution of variance among the 14 q_i 's... 8
- 4 The first ten orthogonal functions of space... 10-12
- 5 Power spectra of \bar{h}' and the first ten orthogonal functions
of time with 43 degrees of freedom... 14-17
- 6 Coherence CH between first ten q_i 's and \bar{h}' as a function of
ordinate number n or period... 19
- 7 Six-day cyclic 100-mb pressure pattern related to \bar{h}' ... 21, 22
- 8 Variations of \bar{h}' and first ten q_i 's from 12/10/56 to 1/15/57... 24
- 9 The 100-mb pressure pattern from 12/28/56 to 1/1/57... 25
- 10 Coherence CH between q_2 and q_5 , q_2 and q_8 , and q_5 and q_8 ... 26

INTRODUCTION

Many studies have been directed toward the common goal of determining the relationship between the tenuous ionosphere and the considerably denser atmosphere below, the stratosphere and troposphere.^{1,2} (See list of references at end of report.) This study examines the relationship between the minimum virtual E -layer height $h'E$ and the 100-mb circulation, as a function of frequency and scale size of the circulation.

DEFINITION OF VARIABLES

$h'E$ is a function of the temperature structure of the atmosphere, solar radiation, atmospheric composition, and recombination rates. $h'E$ represents the lowest region of the ionosphere regularly measured and is essentially the height of the base of the E -layer, usually at about 100 km. $h'E$ at Washington, D. C., was chosen to represent the ionosphere and was taken from the CRPL-F series published by the National Bureau of Standards.³ The height is tabulated in kilometers for every hour during the day when the layer exists. The error of $h'E$ was ± 10 km before January 1956 and ± 2 km afterwards. These large errors mask most variations thought to be of interest, but they were effectively reduced by taking a seven-hour average \bar{h}' of $h'E$ centered around local noon each day. \bar{h}' should be a better daily value of $h'E$ than a single hourly value.

It was decided to examine the relationship between the circulation on a constant pressure surface and \bar{h}' because both are functions of the atmospheric temperature structure. The 100-mb winds at the seven stations shown in figure 1 were chosen to represent the 100-mb circulation,




Figure 1. The seven stations used to describe the 100-mb circulation.

The number of stations was limited by the capacity of the computer for the statistical techniques chosen and the sample size of 2922 days. The station separation was selected so that relatively large- and small-scale sizes of the circulation would be included. The 100-mb winds were supplied by the National Weather Records Center for the eight-year period from March 1951 through February 1959.

The wind directions were given to 16 points of the compass, and the speeds were given in meters per second. The daily u - and v - components of the winds were computed for each station. The u -component is positive toward the east and the v -component is positive to the north.

The u - and v -components and \bar{h}' were filtered to remove the annual cycle in the means. The filter response given in figure 2 shows that the amplitude reduction of the annual cycle was 0.05 while the amplitude of all variations with periods less than 100 days was unaffected. The sample size for the filtered data was 2722 days.




Figure 2. Response curve of the mathematical filter used to filter the data.

CIRCULATION INDICES

It is possible that \bar{h}' is related to certain scale sizes of the 100-mb circulation at certain frequencies. A factor analysis technique developed in the Department of Meteorology at the Massachusetts Institute of Technology⁴ was used to generate circulation indices which represent various scale sizes. The technique produces what has been labeled "empirical orthogonal functions." The technique was used to generate 14 uncorrelated time-varying weighting factors q_i ($1 \leq i \leq 14$) of 14 corresponding orthogonal circulation patterns y_i from the correlated daily values of the seven pairs of filtered u - and v - components. This technique factorized the total variance of the correlated u - and v - components into a minimum number of orthogonal variables q_i 's which contain a certain per cent of the total information. The q_i 's and y_i 's make up the "empirical orthogonal functions." Those q_i 's containing little variance can generally be considered to be noise with respect to application in prediction and need not be correlated with \bar{h}' . Out of the 14 q_i 's generated from the 14 u - and v - components, ten q_i 's having the largest variance were required to contain 90 per cent of the information. The remaining four q_i 's with small variance were eliminated in the correlation analysis. The first ten q_i 's were the circulation indices used in this study. The distribution of the variance among the 14 q_i 's is shown in figure 3.




Figure 3. The distribution of the variance among the 14 q_i 's.

The u - and v - components, the q_t 's, and the y_t 's are related according to the matrix equation:

$$\begin{pmatrix} u_{11} \dots u_{17} v_{11} \dots v_{17} \\ \dots \\ u_{n1} \dots u_{n7} v_{n1} \dots v_{n7} \end{pmatrix} = \begin{pmatrix} q_{11} \dots q_{1,10} \\ \dots \\ q_{n1} \dots q_{n,10} \end{pmatrix} \begin{pmatrix} u y_{11} \dots u y_{17} v y_{11} \dots v y_{17} \\ \dots \\ u y_{10,1} \dots u y_{10,7} v y_{10,1} \dots v y_{10,7} \end{pmatrix} \quad (1)$$

where n is 2722 days. Each row in the $u-v$ matrix is the set of u - and v - components for the seven stations on day t . Each column i in the q -matrix is a function of time t and is orthogonal to the other nine columns. Each row i in the y -matrix is a function of space and is orthogonal to the other nine rows. The u - and v -components for station j on day t are

$$u_{tj} = \sum_{i=1}^{10} q_{ti} u y_{ij} \quad (2a)$$

$$v_{tj} = \sum_{i=1}^{10} q_{ti} v y_{ij} \quad (2b)$$

From equation 2 the direction and magnitude of all winds can be determined with the noise removed. If all q_t 's are zero except when $t = s$, then

$$|\vec{V}| = |q_{ts} (u y_{sj}^2 + v y_{sj}^2)^{\frac{1}{2}}| \quad (3a)$$

$$\tan \theta_j = \frac{v y_{sj}}{u y_{sj}} \quad (3b)$$

When q_s is negative, the wind direction is opposed to the direction given by equation 3b.

Figure 4 shows the first ten orthogonal functions of space y_t 's derived from equations 3a and 3b. The isopleths are relative height contours in feet for q_s equal to 1. It is noticed that, in general, the scale size of the circulation decreases as t increases. This agrees with earlier studies. It is now evident that a correlation between \bar{h}' and the q_t 's, the weighting factors of the y_t 's, is a correlation between \bar{h}' and various scale sizes of the 100-mb circulation.




Figure 4. The first ten orthogonal functions of space. The isopleths are relative height contours of the 100-mb surface when $q_1 = 1$. The wind speeds given at each station are in meters per second.







Figure 4. (Continued)




Figure 4. (Continued)

POWER SPECTRUM

If the relationship between \bar{h}' and the q_t 's is to be examined as a function of frequency or period, as well as a function of scale size, cross-spectrum analysis between \bar{h}' and the q_t 's is required. Cross-spectrum analysis gives the square of the correlation, the coherence CH , as a function of period, and the phase lag τ between correlated frequencies. Two time series can be related at certain frequencies which are not predominant in either time series. An examination of the power spectrum of \bar{h}' and the q_t 's will indicate the relative amplitudes of all frequencies present in the time series.

Figure 5 shows the unnormalized power spectra $U(n)$ of \bar{h}' and the first ten q_t 's as a function of the ordinate number n for n between 0 and 80. The frequency interval between ordinates is equal to 4×10^{-3} cycles per day. Thus the periods range from infinity to 3.3 days. There are 43 degrees of freedom in each estimate of $U(n)$. All power spectra generally show a concentration of energy at low frequencies and a gradual decrease of energy as frequency increases, a common feature of meteorological data. The concentration of variance in the lower order q_t 's should be noted.

No predominant periodicities are present in the power spectra except the spectrum for q_2 . There are two very significant periodicities in q_2 as indicated by the 90 per cent confidence limits.* The two predominant periods are 12.5 and 22.2 days. A power spectrum of q_2 with 20 degrees of freedom, which is not shown, indicates an even greater level of significance at the two periods and places the predominant periods at 12.5 and 21.7 days. The presence of two predominant periodicities in q_2 implies

*The true value of $U(n)$ will lie within the limits given in figure 5 for 90 per cent of the time when there are 43 degrees of freedom.

that y_2 varies every 12.5 and 22 days from a cyclonic to an anticyclonic circulation as q_2 goes from a negative to a positive value. The amplitudes of the two predominant periods are, of course, additive. No pressure pattern motion is implied because y_2 is a fixed pattern of space. If y_2 is part of a wave pattern around the Northern Hemisphere, then y_2 is part of a standing wave pattern. y_2 extends about 3000 km in the east-west direction. If y_2 is one-half a 6000-km wave, then y_2 suggests a wave number of 5. Based on a numerical experiment, Phillips⁵ predicted that the wave number 5 would have an eastward velocity of 1800 km/day. It is interesting to note that the beat between a 12.5 and 21.7 day period is a 29.5 day period, which is almost a lunar cycle.




Figure 5. The power spectra of \bar{h}' and the first ten orthogonal functions of time with 43 degrees of freedom. The frequency interval between the ordinates is 4×10^{-3} cycles/day. The amplitudes of the low frequencies have been reduced by filtering the wind components (see fig. 2).





Figure 5. (Continued)





Figure 5. (Continued)




Figure 5. (Continued)

CORRELATION BETWEEN \bar{h}' AND THE 100-mb CIRCULATION

Figure 6 shows the coherence CH between \bar{h}' and the first ten q_i 's for periods between 250 and 5 days. The CH between two series of eight-digit random numbers having the same degrees of freedom is shown for comparison. Although independence between the q_i 's at all periods has not been established, the CH 's of all ten q_i 's were added to give the total CH , as shown at the bottom as CH_T . If the ten q_i 's are orthogonal at all periods, then CH_T is equivalent to the total reduction in variance. CH_T can be used as a guide in determining the most important periods until the degree of independence between the q_i 's has been established.

There are five predominant peaks in the ten CH functions in figure 6. Two peaks occur with q_2 , one occurs with q_5 , and two occur with q_8 . q_2 is related to \bar{h}' at a period of about 13 or 14 days and at a period of 6.3 days. The shape of the peak regions in CH for q_5 and q_8 are surprisingly similar between the 6.3 and 6.6 day period. q_8 also has a peak in CH at a period of 17.9 days. The probability that the five peak values of CH could happen by chance ranges from 0.165 to 0.035 as shown in figure 6. Although the five CH peaks are not highly significant, an examination of the relationship between \bar{h}' , q_2 , q_5 , and q_8 at a period of 6.3 days was thought to be valuable primarily because the relationship apparently involves the fundamental periodicity of 12.5 days found to be predominant in q_2 . CH_T also indicates that the relationship between \bar{h}' and the 100-mb level is significant at a period of 6.3 days. The remaining part of this paper will be an examination of the relationship between \bar{h}' , q_2 , q_5 , and q_8 at a period of 6.3 days.

The phase lag τ in days for q_2 , q_5 , and q_8 , relative to \bar{h}' for a period of 6.3 days, is 2.4, 4.4, and 6.2 days respectively. The 90 per cent confidence limits of these




Figure 6. The coherence CH between the first ten q_i 's and \bar{h} 's as a function of the ordinate number n or period. The most significant regions of CH have been accented.

lags are ± 0.7 , ± 0.8 , and ± 0.7 days respectively. * None of the 90 per cent limits overlap. q_8 can be assumed to be in phase with \bar{h}' because the lag of 6.2 days is essentially equal to the period of 6.3 days. Assuming that the phase lags are 2, 4, and 0 days for q_2 , q_6 , and q_8 , respectively, and that the relative amplitudes at a period of 6.3 days are given by the power spectra, the cyclical pressure pattern related to \bar{h}' can be constructed.

Figure 7 gives the cyclic pressure pattern related to \bar{h}' as specified by q_2 , q_6 , and q_8 for an even six-day period instead of a 6.3-day period. The isopleths in figure 7 are relative heights of the 100-mb surface for the relative wind speeds shown. On Day 1 (4) a trough (ridge) extends in the east-west direction. On Day 2 (5) the trough (ridge) begins to tilt in a NE-SW direction. On Day 3 (6) the southern part of the trough (ridge) moves eastward, while the northern position remains about stationary, but the general NE-SW tilt is conserved. The maximum (minimum) in \bar{h}' occurs on Day 3 (6). The pressure trough on Day 3, when h' peaks, is quite similar to a wave on a frontal surface. Apparently \bar{h}' is associated with NE-SW oriented ridges and troughs on the 100-mb surface, which occur about every six days. The maximum of \bar{h}' occurs when q_2 is negative and increasing; that is, when the cyclonic flow is decreasing and being replaced by an anticyclonic flow. The phase lag between \bar{h}' and q_2 at a period of about 13 days is about 10 days, which indicates that the maximum of \bar{h}' occurs when q_2 is about zero and decreasing in magnitude.

*The true value of τ will lie within the stated limits 90 per cent of the time when there are 43 degrees of freedom.







Figure 7. The six-day cyclic 100-mb pressure pattern that is related to \bar{h}' . The isopleths are relative height contours of the 100-mb surface.





Figure 7. (Continued)







Figure 8. Variations of \bar{h}' and the first ten q_i 's from 10 December 1956 to 15 January 1957.




100 - mb Q - CIRCULATION
DECEMBER 28, 1956




100 - mb Q - CIRCULATION
DECEMBER 29, 1956



100 - mb Q - CIRCULATION
DECEMBER 30, 1956



100 - mb Q - CIRCULATION
DECEMBER 31, 1956



100 - mb Q - CIRCULATION
JANUARY 1, 1957

Figure 9. The 100-mb pressure pattern from 28 December 1956 to 1 January 1957 according to the first ten q_i 's given in figure 8. The isopleths are relative height contours for the 100-mb surface.

COHERENCE BETWEEN THE CIRCULATION INDICES

To determine the use of CH_T as an indicator of the total reduction in variance, CH between q_2 and q_5 , q_2 and q_8 , and q_5 and q_8 were determined. These CH 's are shown in figure 10. It should be noted that relatively large peaks occur in CH at a period of about 6.3 days, particularly between q_2 and q_8 and between q_5 and q_8 . It should be noted that neither q_2 , q_5 , nor q_8 has a predominant periodicity at a period of 6.3 days. The peak between q_5 and q_8 has a p -value of 0.052 and is more significant than the other peaks. The similarity in the shape of the CH curves of $\bar{h}'-q_5$ and $\bar{h}'-q_8$ at a period of about six days probably resulted from the correlation between q_5 and q_8 at about a six-day period. Thus, it appears that CH_T can be used only to indicate the most important regions where \bar{h}' and the 100-mb circulation are related.




Figure 10. The coherence CH between q_2 and q_5 , q_2 and q_8 , and q_5 and q_8 .

CONCLUSIONS

From the standpoint of statistical significance, the most important result of this study is the detection of two predominant periodicities in one of the ten orthogonal indices used to describe the 100-mb circulation. These predominant periods occur in q_2 and are 22 and 12.5 days. Apparently \bar{h}' and the 100-mb circulation are only related at periods of about 18, 13, and 6 days. The correlation between \bar{h}' and q_2 at an 18-day period was not examined in this study. The mechanisms producing the 22 and the 12.5 day period in q_2 may be different because \bar{h}' is not related to the 100-mb circulation at the 22-day period while it is at the 12.5-day period. It is thought that the statistically deduced pressure pattern related to \bar{h}' at a period of about 6.3 days is significant although the significance of the CH peaks at a period of 6.3 days is not high because the 6.3-day pattern is apparently controlled by the predominant 12.5-day period.

RECOMMENDATION

Correlate other ionospheric parameters with the circulation over larger regions of the world and at other pressure levels in the stratosphere or troposphere. Examine the physics of the relationship between $h'E$ and the stratospheric circulation particularly with respect to the preference for $h'E$ to be related with certain circulation patterns which occur at certain periods.

REFERENCES

1. Craig, R. A., "Surface-Pressure Variations Following Geomagnetically Disturbed and Geomagnetically Quiet Days," Journal of Meteorology, v. 9, p. 126-138, April 1952
2. Shapiro, R., "Further Evidence of a Solar-Weather Effect," Journal of Meteorology, v. 13, p. 335-340, August 1956
3. National Bureau of Standards. Central Radio Propagation Laboratory, Series F, Part A: Ionospheric Data
4. Massachusetts Institute of Technology. Meteorology Department Contract AF19(604)-1566; Scientific Report 1, Empirical Orthogonal Functions and Statistical Weather Prediction, by E. N. Lorenz, December 1956
5. Phillips, N. A., "General Circulation of the Atmosphere: A Numerical Experiment," p. 18-25 in Conference on the Application of Numerical Integration Techniques to the Problem of the General Circulation. Dynamics of Climate; Proceedings. Held October 26-28, 1955, Pergamon Press, [1960]

INITIAL DISTRIBUTION LIST (continued)

Redstone Scientific Information
 Center
 Army Electronic R&D Laboratory
 SELRA/SL-XC SELRA/SL-ADT
 SELRA/SR
 Electronic Warfare Division
 Picatinny Arsenal
 Army Research Office (Durham)
 Internal Research Division
 Electronic Development Activity
 White Sands
 Army Electronic Research Unit
 Army Airborne, Ft. Bragg
 Deputy Chief of Staff, Development,
 USAF AFRST-EL/CS
 Deputy Chief of Staff, Operations,
 USAF AFOCC-C11
 Air Defense Command
 ADOAC-DL ADOOA
 Air Research & Development
 Command SCSE
 Air University, Library AUL3T-5028
 Alaskan Air Command
 Electronics (2)
 Strategic Air Command, OAST
 Air Force Cambridge Research
 Center AFRD, ERD
 Air Force Missile Test Center
 MU-135
 Rome Air Development Center
 RAALD
 Wright Air Development Division
 ASAPRD-Dist ASNVEG
 ASROO-3 (2)
 Air Proving Ground Center
 PGAPI
 1st Strategic Aerospace Division
 DOEO
 Norton Air Force Base
 Federal Aviation Agency
 Navigation Branch
 NASA, Langley Research Center (3)

Executive Secretary, Armed Forces-NRC
 Vision Committee
 Air Weather Service, Scott AFB
 National Security Agency
 C121 (2) R31
 National Bureau of Standards,
 Boulder Library
 Central Radio Propagation Lab
 Federal Communications Commission
 U. S. Weather Bureau
 Washington, D. C. (2)
 Norman, Okla.
 National Science Foundation
 Defense Com. Agency
 University of California
 Engineering Dept.
 Stanford University
 Electronics Lab.
 University of Alaska, Geophysical
 Institute
 University of California, Space
 Science Lab., Berkeley
 California Institute of Technology,
 Chief, Jet Propulsion Lab.
 University of Miami
 Marine Laboratory (3)
 Cornell University, Library
 Dartmouth College, Thayer School
 of Engineering
 Georgia Institute of Technology,
 Engineering Experiment Station
 Harvard Observatory, Dr. Menzel (2)
 New York University
 Institute of Mathematical Sciences
 Dept of Meteorology & Oceanography
 The Ohio State University
 Rocket Research Lab
 Pennsylvania State University
 Ordnance Research Lab
 University of Texas, Electrical
 Engineering Research Lab
 University of Michigan, Electrical
 Engineering Department
 Lamont Geological Observatory

<p>Navy Electronics Laboratory Report 1155</p> <p>CORRELATION BETWEEN THE MINIMUM VIRTUAL F-LAYER HEIGHT $h'f$ AND THE 100-MB CIRCULATION, by V. R. Noonkester, 28p., 22 January 1963.</p> <p>UNCLASSIFIED</p> <p>A study was conducted to examine the correlation between the minimum virtual F-layer height $h'f$ at Washington, D. C. and the stratospheric circulation over the Eastern U. S. as a function of frequency and scale size of the circulation. The minimum virtual F-layer height was found to be correlated with a particular type of large-scale circulation at the 100-mb level which has a period of 6, 3 days. One of the ten circulation indices used to describe the 100-mb circulation was found to have predominant periods of 22 and 12 days.</p> <p>1. Atmosphere - Temperature-Electromagnetic effects 2. Ionosphere - Electromagnetic effects 1. Noonkester, V. R.</p> <p>SR 008 01 01, Task 7039 (NEL M2-3) SF 011 05 11, Task 0558 (NEL M2-5)</p> <p>This card is UNCLASSIFIED.</p>	<p>Navy Electronics Laboratory Report 1155</p> <p>CORRELATION BETWEEN THE MINIMUM VIRTUAL F-LAYER HEIGHT $h'f$ AND THE 100-MB CIRCULATION, by V. R. Noonkester, 28p., 22 January 1963.</p> <p>UNCLASSIFIED</p> <p>A study was conducted to examine the correlation between the minimum virtual F-layer height $h'f$ at Washington, D. C. and the stratospheric circulation over the Eastern U. S. as a function of frequency and scale size of the circulation. The minimum virtual F-layer height was found to be correlated with a particular type of large-scale circulation at the 100-mb level which has a period of 6, 3 days. One of the ten circulation indices used to describe the 100-mb circulation was found to have predominant periods of 22 and 12 days.</p> <p>1. Atmosphere - Temperature-Electromagnetic effects 2. Ionosphere - Electromagnetic effects 1. Noonkester, V. R.</p> <p>SR 008 01 01, Task 7039 (NEL M2-3) SF 011 05 11, Task 0558 (NEL M2-5)</p> <p>This card is UNCLASSIFIED.</p>
<p>Navy Electronics Laboratory Report 1155</p> <p>CORRELATION BETWEEN THE MINIMUM VIRTUAL F-LAYER HEIGHT $h'f$ AND THE 100-MB CIRCULATION, by V. R. Noonkester, 28p., 22 January 1963.</p> <p>UNCLASSIFIED</p> <p>A study was conducted to examine the correlation between the minimum virtual F-layer height $h'f$ at Washington, D. C. and the stratospheric circulation over the Eastern U. S. as a function of frequency and scale size of the circulation. The minimum virtual F-layer height was found to be correlated with a particular type of large-scale circulation at the 100-mb level which has a period of 6, 3 days. One of the ten circulation indices used to describe the 100-mb circulation was found to have predominant periods of 22 and 12 days.</p> <p>1. Atmosphere - Temperature-Electromagnetic effects 2. Ionosphere - Electromagnetic effects 1. Noonkester, V. R.</p> <p>SR 008 01 01, Task 7039 (NEL M2-3) SF 011 05 11, Task 0558 (NEL M2-5)</p> <p>This card is UNCLASSIFIED.</p>	<p>Navy Electronics Laboratory Report 1155</p> <p>CORRELATION BETWEEN THE MINIMUM VIRTUAL F-LAYER HEIGHT $h'f$ AND THE 100-MB CIRCULATION, by V. R. Noonkester, 28p., 22 January 1963.</p> <p>UNCLASSIFIED</p> <p>A study was conducted to examine the correlation between the minimum virtual F-layer height $h'f$ at Washington, D. C. and the stratospheric circulation over the Eastern U. S. as a function of frequency and scale size of the circulation. The minimum virtual F-layer height was found to be correlated with a particular type of large-scale circulation at the 100-mb level which has a period of 6, 3 days. One of the ten circulation indices used to describe the 100-mb circulation was found to have predominant periods of 22 and 12 days.</p> <p>1. Atmosphere - Temperature-Electromagnetic effects 2. Ionosphere - Electromagnetic effects 1. Noonkester, V. R.</p> <p>SR 008 01 01, Task 7039 (NEL M2-3) SF 011 05 11, Task 0558 (NEL M2-5)</p> <p>This card is UNCLASSIFIED.</p>

INITIAL DISTRIBUTION LIST
(Continued on inside back cover)

Bureau of Ships

Code 320 Code 690 (2)
Code 360 (3) Code 210L (3)
Code 670 Code 673
Code 686 Code 687A (2)
Code 684

Bureau of Naval Weapons

DLI-3 DLI-31
RAAV-6 RAAV-321
RUDC-11 RUDC-2
R-56 FASS

Chief of Naval Personnel

Code 11B

Chief of Naval Operations

Op-07T Op-03EG
Op-94 Op-94G43

Chief of Naval Research

Code 418 Code 427 Code 461

Commander in Chief, Pac Flt

Commander in Chief, Lant Flt

**Commander Operational Test &
Eval. For, Lant Flt**

**Deputy Commander Operational
Test & Eval. For, Pac**

**Commander Training Command,
Pac Flt**

**Commander Submarine Development
Group TWO**

**Commander, Amphibious Force,
Pac Flt (2)**

Commander Service For, Lant Flt

**Naval Air Development Center,
Library**

Naval Missile Center

Tech Library Code 5320
Code 3250

Naval Air Test Center

NANEP WST-424

**Naval Ordnance Laboratory,
Library**

**Naval Ordnance Test Station,
Pasadena**

**Naval Ordnance Test Station,
China Lake Code 753**

Naval Weapons Laboratory

Naval Radiological Defense Lab

David Taylor Model Basin, Library

**Naval Engineering Experiment
Station, Library**

**Navy Underwater Sound Laboratory
Code 1450 (3)**

Naval Research Laboratory

Code 5400 Code 2027

Code 5250 Code 5330

**Navy Underwater Sound Reference
Laboratory, Library**

**Naval Ordnance Laboratory, Corona
Beach Jumper Unit I**

Beach Jumper Unit II

Air Development Squadron ONE (VX-1)

**Office of Naval Research, Pasadena
Boston**

**Naval Submarine Base, New London
Electronics Officer**

Navy Hydrographic Office (2)

**Naval Postgraduate School,
Library (2)**

**U. S. Naval Personnel Research
Field Activity**

**U. S. Navy Weather Research
Facility**

**Navy Representative, Project
LINCOLN, MIT**

**Defense Atomic Support Agency
DOD, Research & Engineering,
Tech. Library**

**Assistant Chief of Staff, G-2,
US Army**

**Aberdeen Proving Ground
Tech. Library (2)**

Ballistics Research Lab

Hdqtrs: AMSTE-EL (2)

**Army Electronic Proving Ground
Tech. Library**